

## CTスキャンは大動脈弁置換術後の弁尖の動きの低下を可視化する(Abstract 17-LB-16234)

抗血小板薬ではなく経口抗凝固薬は無症候性の弁尖血栓症を減少させる

Oral anticoagulants, but not anti-platelet therapy associated with reduced subclinical leaflet thrombosis

大動脈弁置換術を施行された患者の約12%が、弁の動きの低下する無症候性の弁尖血栓症を発症したとの観察研究が、American College of Cardiology's 66<sup>th</sup> Annual Scientific Session で発表され、同時にLancetに掲載された。CTスキャンの解析の結果、経カテーテル大動脈弁置換術(TAVR)施行患者の13.6%および外科的大動脈弁置換術(SAVR)施行患者の3.8%において、無症候性弁尖血栓が認められ、全体では12.1%に認められた。血栓はTAVR患者においてSAVR患者よりも有意に高率に認められたが、この差はSAVR施行患者の方が若年であり全体的な健康状態が良好であることによると考えられる。

### Full Text

About 12 percent of patients undergoing aortic valve replacement developed non-symptomatic subclinical leaflet thrombosis that reduced the motion of the valves, according to an observational study presented at the American College of Cardiology's 66<sup>th</sup> Annual Scientific Session.

The study is the largest to date investigating thrombosis as a potential cause of reduced valve motion after aortic valve replacement. It confirms a previous, smaller study that suggested blood clots that are detectable with computed tomography (CT) scans but not with more commonly-used echocardiogram scans can develop around the valve and constrain the valve's motion. In the new study, the CT-detected valve-associated clots were found to increase the risk of transient ischemic attacks, also called "mini-strokes," but were not associated with an increased risk of death, heart attack or stroke. Anticoagulant therapy, but not anti-platelet therapy, was associated with a significantly lower risk of developing valve-associated clots.

"This phenomenon of subclinical leaflet thrombosis can be missed if you just use transthoracic echocardiogram," said Raj Makkar, MD, associate director of Cedars-Sinai Heart Institute in Los Angeles, and the study's lead author. "Based on our study, CT is clearly a more sensitive and appropriate technique to actually make a diagnosis of subclinical leaflet thrombosis. This suggests clinicians might want to have a lower threshold to do a CT scan if there is suspicion of reduced motion in the valve, such as from slightly elevated mean gradients on echocardiogram."

CT scans expose patients to more radiation than echocardiogram alone.

The researchers analyzed CT scans and other health records from 850 patients enrolled in two single-center medical registries, known as RESOLVE (which includes patients treated at Cedars-Sinai Heart Institute) and SAVORY (which includes patients treated at Rigshospitalet hospital in Copenhagen, Denmark). The patients had undergone CT scans an average of three months after transcatheter aortic valve replacement (TAVR) or surgical aortic valve replacement (SAVR), two procedures used to replace a patient's faulty aortic valve with an artificial valve. In TAVR, clinicians thread the new valve to the heart through a catheter in the groin or chest; this procedure is generally used for patients at intermediate to high cardiovascular risk, which typically includes older patients and those who have multiple health problems. SAVR is an open-heart procedure used in lower-risk patients.

A total of 720 patients in the two registries had undergone TAVR and 130 had undergone SAVR. Analysis of CT scans revealed subclinical leaflet thrombosis in 13.6 percent of TAVR patients and 3.8 percent of SAVR patients, for an overall rate of 12.1 percent among all patients combined. Although thrombosis was observed in a significantly greater proportion of TAVR patients than SAVR patients, this difference may be attributable to the younger average age and better overall health of patients undergoing SAVR, Makkar said.

The results also showed that subclinical leaflet thrombosis was significantly more common in patients on antiplatelet therapy (typically aspirin plus a P2Y12 inhibitor) compared to those taking anticoagulants. A total of 14.8 percent of patients on antiplatelet therapy had thromboses compared to 4 percent among patients taking warfarin and 3 percent among patients taking non-vitamin K antagonist anticoagulants, or NOACs. There was no significant difference in risk observed among those taking warfarin versus NOACs.

"We need to further study whether routine anticoagulation may be useful for this patient population," Makkar said. "Dual antiplatelet therapy was not effective in preventing and treating subclinical leaflet thrombosis, and it does have a small risk of bleeding, particularly in older patients. There is an impetus to study the risks and benefits of dual antiplatelet therapy further in randomized clinical trials."

An analysis of a small group of patients (58) who underwent a second CT scan showed subclinical leaflet thrombosis resolved over time in the vast majority of patients who were started on oral anticoagulant therapy after the first CT scan and that thromboses resolved in only a small portion of patients who were not started on anticoagulants.

The study also showed that subclinical leaflet thromboses detected with CT scans were reflected in significantly higher mean gradients, a measure used to assess functioning of the aortic valve using echocardiogram. However, Makkar said this difference was not large enough to enable clinicians to diagnose subclinical leaflet thrombosis using echocardiogram alone.

As an observational study, the results do not directly assess cause and effect. Makkar said ongoing randomized clinical trials that include CT scans as part of the protocol should help to further elucidate the factors that contribute to subclinical leaflet thrombosis after aortic valve replacement.

"Our study findings can help optimize the use of different blood thinning medications in patients undergoing aortic valve replacement, which might potentially result in further improvements in valve hemodynamics and clinical outcomes," Makkar said.

This study was simultaneously published online in The Lancet at the time of presentation.

## ACC2017特集

[News01]

エボロクマブは心血管イベントを有意に低下させた関連付ける

[News02]

自己拡張型経カテーテルの大動脈弁置換術(TAVR)は中等度リスクの患者に適している

[News03]

NSAIDに加えてミソプロストールを服用することにより心血管系リスクが低下する

[News04]

手首装着型心拍計は胸部装着型よりも正確さに欠ける

[News05]

ホルモン補充療法は死亡率が低いことと関連がある

[News06]

MI後の睡眠時無呼吸スクリーニングに最適な時期が調査された

[News07]

運動歴は乳がん後の心疾患予防に役立つ

[News08]

うつ病はMIまたは狭心症後の死亡リスクを倍増させる

[News09]

MI直後にすべての閉塞動脈を治療することによる有益性

[News10]

リバーロキサバンはアスピリンに比べVTE再発を軽減する

[News11]

CTスキャンは大動脈弁置換術後の弁尖の動きの低下を可視化する

[News12]

BococizumabによるPCSK9阻害による結果は様々である

[News13]

TAVRは微小出血および神経学的障害と関連がある

[News14]

スタチン服用患者においてエボロクマブは認知機能に影響しない

[News15]

左心耳閉鎖術は脳卒中リスクを低下させる

[News16]

ペースメーカープログラムは意識消失発作を減少させる

[News17]

心房細動患者においてジゴキシンにより死亡リスクは上昇する

[News18]

新規抗凝固薬は心房細動に対するアブレーション中の大出血を軽減する

[News19]

血液検査により非心臓手術後の心損傷が検出できる

[News20]

ウェブベースのカウンセリングは血圧を低下させる