

積極的なモニタリングはAFibの診断率を3倍に上昇させる (Abstract LB-18937-ACC)

mSToPS: 自己装着胸部パッチは一般的な不整脈を従来の方法よりもより迅速に捕捉できる
mSToPS: Self-applied chest patch catches common irregular heartbeat more quickly than usual care

心房細動(AFib)が高リスクの人々において、心調律を記録する着用型自己装着胸部パッチは、従来の方法に頼るよりも不整脈の検出に優れより迅速な治療を促進する可能性があるとの1年間のデータが、American College of Cardiology's 67th Annual Scientific Session で発表された。mHealth Screening to Prevent Strokes (mSToPS) トライアルの対象患者は心電図を持続的に記録するパッチを装着し、その結果研究者らは連続14日間のデータを得ることができた。AFibは、装着型パッチ群の6.3% およびコントロール群の2.3% において新たに診断された。パッチを用いた積極的なモニタリングにより、抗凝固薬開始率が有意に高まった(5.4% vs. 3.4%)。

Full Text

For people at heightened risk for atrial fibrillation (AFib), wearing a self-adhering chest patch that records heart patterns may better detect the condition and facilitate more timely treatment than relying on usual care, according to one-year data presented at the American College of Cardiology's 67th Annual Scientific Session.

The study, called the mHealth Screening to Prevent Strokes (mSToPS) trial, is one of the first completely digital, nationwide, direct-to-participant and site-less clinical research programs, according to researchers. Patients consented and enrolled exclusively via a web-based platform to undergo active monitoring at home using the iRhythm Zio patch. The patch—about the size of a large Band-Aid – continuously records an electrocardiogram (ECG), the gold standard for detecting AFib. But instead of only providing a brief snapshot of someone's heart rhythm – which is often the case with an ECG in the clinic – this technology allowed researchers to collect 14 consecutive days of data capturing the electrical activity of an individual's heart.

"At one year, people who wore the chest monitor had nearly three times the likelihood of being diagnosed with AFib appropriately. A significantly higher proportion of these patients were started on anticoagulant therapy to lower their stroke risk compared with those who received usual care," said Steven R. Steinhubl, MD, director of digital medicine, Scripps Translational Science Institute, La Jolla, California and lead author of the study. "The data also provide a first glimpse into how this type of ECG screening might influence heart care utilization."

People with AFib have a fivefold greater chance of having a stroke than the general population. New estimates suggest that 37 percent of adults over 55 years of age will develop AFib. But Steinhubl said as many as 1 out of 3 patients may be undiagnosed and, therefore, remain susceptible to stroke.

"A stroke is a very devastating cardiovascular complication of AFib; it can be tragic for patients and costly for payers," Steinhubl said. "If we identify and get these people on a blood thinner, we can reduce their risk of stroke by about 70 percent, which is powerful."

The mSToPS trial enrolled and monitored a total of 1,732 eligible Aetna patients with no heart rhythm issues but who were at moderate risk for AFib (median CHA2DS2-VASc score of 3) and matched them (2:1) to 3,646 observational controls of similar age, sex and risk profiles who received usual care only; all were included in the analysis. The average patient age was 73.7 years.

Once enrolled, participants in the active monitoring group received the wearable chest patch by mail and were given clear written and video instructions on how to use and place the wearable patch. Participants wore it for an average of 12 days and then returned it with a prepaid shipping envelope. The data collected from the patch, currently only U.S. Food and Drug Administration-approved for clinical use, are first analyzed with an algorithm that reads the ECG and preliminarily identifies all rhythm abnormalities, including AFib. In the trial analyses, a clinician and a clinical events committee read all potential diagnoses of AFib in a blinded fashion and either agreed or disagreed. All participants received their ECG study report. Researchers also collected and analyzed use of AFib-related medications, doctor and emergency department (ED) visits, and cases of blood clots and stroke.

The primary endpoint was the incidence of AFib at one year. AFib was newly diagnosed in 6.3 percent of those wearing the patch and 2.3 percent of controls. Compared with controls, participants who wore the chest patch had significantly more primary care physician visits (78.7 vs. 75 percent,) and cardiology outpatient visits (31.6 vs. 23.6 percent), but no difference in ED visits or hospitalizations. Relative to controls, active monitoring with the patch was associated with significantly greater initiation of anticoagulation (5.4 vs. 3.4 percent) and small but statistically significant increases in antiarrhythmic therapy (0.8 vs. 0.3 percent) and pacemaker or implantable cardioverter-defibrillator placement (0.7 vs. zero percent). Researchers were unable to show that the intervention had an impact on clinical outcomes, including blood clots or stroke, given the short follow-up (eight months on average). Study participants will be followed for another two years to track these events.

Still, Steinhubl said the findings demonstrate a feasible, scalable and clinically valuable way of screening for undiagnosed AFib in an at-risk, nationwide population.

"The quality of data collected through the patch is as good as what we see clinically," he said. "What was fascinating is that for the people with AFib, the burden of AFib (the amount of time a person was in AFib relative to their entire monitoring time) was quite low – approximately 1 percent of the time on average." This finding that suggests the prevalence of undiagnosed AFib is likely higher than clinicians might suspect, since it can be easy to miss, which could have important implications for how we should ideally screen for the condition.

"In most cases, the only currently recommended method is to feel someone's pulse or check an ECG for 30 seconds during a routine doctor's visit," Steinhubl said. "Based on our data, individuals have very short episodes of AFib that would make it very difficult to catch in the way we routinely look for it today."

Increasing age is by far the biggest risk factor for AFib, but heart failure, prior stroke and sleep apnea are all additional important risk factors. Steinhubl stressed that one of the most unique and valuable aspects of the study is how it was designed and conducted, including an emphasis on being more participant-centric. Using digital technology, the study was brought directly to the participants, even participants who were 74 years old on average.

"Currently, most people who are able to take part in clinical research live close to academic medical centers," he said. "In a way, mSToPS is proof of principle that site-less trials are possible and might open up clinical research to anybody, anywhere who is interested in taking part."

Researchers will continue to follow participants for another two years to assess any differences in reports of strokes or other systemic thromboemboli.

This study was funded, in part, by Janssen Pharmaceuticals. Other sponsors include the National Institutes of Health and the Qualcomm Foundation.

ACC2018特集

[News01]
気候変動は心筋梗塞リスクを上昇させる可能性がある

[News02]
炎症性腸疾患はMIリスクを上昇させる

[News03]
前向きな態度は狭心症患者の転帰を改善する

[News04]
アリロクマブは急性冠症候群後の心血管イベントを軽減する

[News05]
着用型自動除細動器は全死亡を減らすが突然死には影響しない

[News06]
心不全患者にとってインフルエンザワクチンは有益である

[News07]
音楽は運動負荷試験中の運動時間を増加させる

[News08]
がん治療は心不全リスクを上昇させる

[News09]
遺伝子型解析はPCI後の薬物選択において有益である

[News10]
3種の低用量内服は高血圧管理に成功した

[News11]
ACSにおけるスタチンのローディングドーズ投与は臨床イベントリスクを減少させない

[News12]
卵円孔閉鎖患者においてデバイスが転帰を改善する

[News13]
ダビガトランは非心臓手術後の心筋障害を軽減する

[News14]
短期抗血小板薬2剤併用療法はMIリスクを上昇させる

[News15]
薬剤が第Ⅹa因子阻害効果をリバーズする

[News16]
Canakinumabは糖尿病への進行を予防しない

[News17]
MI後のチカグレロル使用の安全性はクロピドグレルと同等である

[News18]
積極的なモニタリングはAFibの診断率を3倍に上昇させる

[News19]
化学療法による心毒性の軽減

[News20]
カルベジロールは乳がん女性の心臓を保護する