

メトトレキサートは心血管イベントを減少させない (Abstract LBCT-17778)

CIRT:低用量のメトトレキサートは二次予防において炎症または心血管イベントを減少させなかった

CIRT: Low dose methotrexate fails to reduce inflammation or cardiovascular events in secondary prevention

炎症軽減による心疾患および脳卒中リスク軽減と言えば、待望のCIRT試験の結果、リスクを有する患者において適切な炎症経路を標的とすることが重要であることが示された。昨年、CANTOS試験において、インターロイキン-1 β 阻害薬カナキヌマブが特定の炎症経路を標的とし、その後心梗塞および心血管死亡率を低下させることが示された。対照的に、CIRT試験の結果、低用量メトトレキサートは、この炎症経路は阻害せずまた主要な心血管イベント率も低下させないことが示された。この結果は、American Heart Association Scientific Sessions 2018で発表され、*New England Journal of Medicine*に掲載された。

Full Text

When it comes to reducing inflammation to decrease the risk of heart disease and stroke, results from the much-anticipated Cardiovascular Inflammation Reduction Trial (CIRT) indicate that targeting the right inflammatory pathways in at-risk patients is crucial. Last year, the Canakinumab Anti-inflammatory Thrombosis Outcomes Study (CANTOS) showed that the interleukin-1 β inhibitor canakinumab both targeted a specific inflammatory pathway and consequently lowered rates of myocardial infarction (MI) and cardiovascular death. By contrast, the findings from CIRT showed that low-dose methotrexate neither inhibited that same inflammatory pathway nor reduced major adverse cardiovascular event rates.

These results were presented by Paul Ridker, MD, director of the Center for Cardiovascular Disease Prevention at Brigham and Women's Hospital, during the American Heart Association Scientific Sessions 2018, and published simultaneously in *The New England Journal of Medicine*.

"The contrasting results between these two contemporary clinical trials demonstrate the importance of considering the mechanistic diversity of inflammatory pathways and of approaches to their inhibition," said Ridker. "Understanding these differences will be crucial for future studies targeting inflammation in atherosclerosis."

Prior to CIRT, observational studies had suggested that low-dose methotrexate, an inexpensive and effective drug widely used to treat rheumatoid arthritis and other inflammatory diseases, might reduce rates of cardiovascular events. The federally-funded CIRT was designed to rigorously test whether low-dose methotrexate could effectively reduce risk of major adverse cardiovascular events - that is, MI, stroke and cardiovascular death.

In parallel, Ridker and colleagues also designed and conducted CANTOS, sponsored by Novartis, to test the same outcomes for canakinumab, a drug that specifically targets interleukin-1 β . Interleukin-1 is a pro-inflammatory cytokine that, if over-produced, results in increased inflammation throughout the body as well as elevated levels of interleukin-6 and high sensitivity C-reactive protein (hsCRP), two critical biomarkers of inflammation.

CIRT and CANTOS were both randomized, double-blind, placebo-controlled trials, and both enrolled stable but high-risk atherosclerosis patients. CANTOS, however, was designed to include only patients with persistently elevated hsCRP levels. CIRT did not employ this criterion, and the average hsCRP level for the population was well within the normal range. CIRT enrolled 4,786 North American patients with prior heart attack or multi-vessel coronary disease who additionally had either type 2 diabetes or a metabolic syndrome. The trial stopped after a median follow-up of 2.3 years.

Unlike canakinumab as used in CANTOS, low-dose methotrexate as used in CIRT did not reduce the inflammatory pathway leading from interleukin-1 to interleukin-6 and on to hsCRP. Concordantly, and in contrast to canakinumab, low-dose methotrexate did not lower cardiovascular event rates compared to placebo. The team reports that 201 patients taking methotrexate suffered a major cardiovascular event compared to 207 patients taking the placebo. Yet, methotrexate was associated with elevations of liver enzymes, reductions in leucocytes and hematocrit, and a higher incidence of non-basal cell skin cancers.

"The results from CIRT and CANTOS, when considered together, tell us something critically important: Not all inflammation is the same, and not all drugs that target inflammation are the same," said Ridker. "While it is disappointing that an inexpensive drug like methotrexate did not have the effects we previously saw in CANTOS, the results from CIRT shed crucial light on the underlying biology that connects inflammation with cardiovascular disease. The divergent trial results provide a clear roadmap to guide our efforts going forward."

Despite its wide clinical use, the biological mechanisms underlying the anti-inflammatory effects of methotrexate in rheumatoid arthritis and other inflammatory conditions remains poorly understood. Drugs such as colchicine and oral NLRP3 inhibitors that may intersect with the interleukin 1 to interleukin-6 to CRP pathway are currently under investigation or in development.

"CANTOS and CIRT provide the cardiovascular community proof-of-principle that specific targeting of the interleukin-1 to interleukin-6 pathway of innate immunity is crucial for preventing atherothrombotic events. The research goal now and the clinical need of our patients is to find inexpensive and widely applicable agents that can safely target this pathway," Ridker said.

Ridker and co-authors received investigator-initiated grants from the National Heart Lung and Blood Institute to conduct CIRT. They also received an investigator-initiated grant from Novartis to conduct CANTOS. Ridker is listed as a co-inventor on patents held by Brigham and Women's Hospital that relate to the use of inflammatory biomarkers in cardiovascular disease and diabetes that have been licensed to AstraZeneca and Siemens, and has served as a consultant to Novartis and Inflazome. Co-authors have received compensation for their site efforts enrolling participants into the trial.

This research was supported by the NIH/National Heart Lung and Blood Institute and Novartis.

Cardiology特集

AHA2018 (第91回米国心臓病協会)

トピックス一覧

[News01]

高用量EPAによる心血管疾患予防

[News02]

魚油およびビタミンDの経年による予防効果

[News03]

糖尿病治療薬は心不全を予防する

[News04]

心疾患を有する糖尿病患者においてバイパス手術は血管形成術よりも優れている

[News05]

意思決定支援ツールが心房細動の管理を改善する

[News06]

糖尿病治療薬は心臓の構造を改善する

[News07]

アンジオテンシン受容体ネプリライシン阻害薬はACE阻害薬より優れている

[News08]

メトトレキサートは心血管イベントを減少させない

[News09]

エゼチミブは一次予防目的の標準治療として最良である

[News10]

PTSDは心停止後のリスクを上昇させる

[News11]

冠動脈石灰化は冠動脈リスクの優れた予測因子である

[News12]

慢性的な騒音への曝露は心血管リスクを上昇させる