

鉛への曝露は長期に渡り認知機能に対し影響する (Abstract# SSJ16-06)

小児期の鉛への曝露は長期の認知機能および行動上の問題を引き起こす
Childhood lead exposure contributes to long-term cognitive and behavioral problems

機能的磁気共鳴画像 (fMRI) にて脳機能を評価した結果、小児期に鉛に曝露された成人は認知機能および行動上の長期の影響を有し永続的な脳傷害を患うと2009年 Radiological Society of North America学会 (RSNA 2009) で発表された。このスタディには血液中の鉛レベルが5~37 (平均14) $\mu\text{g/dL}$ の成人33人 (女性14人および男性19人、平均21歳) が組み入れられた。病歴上IQ低下、少年非行および犯罪による逮捕が認められた。全ての患者が、注意力、意思決定および衝動コントロールを司る脳の実行機能を計測する二つの課題を行っている最中のfMRIを施行された。画像検査の結果、抑制を要する課題を完了するために、血中の鉛レベルの上昇している者は前頭葉および頭頂葉の余分な領域の活動性を必要とすることが示され、抑制を司る脳領域が鉛によりダメージを受けていることが示唆された。注意力を検査するようにデザインされた課題を行っている最中の画像検査の結果、鉛レベルが高いと頭頂葉領域および他の脳領域の活性が低下することが示された。これらの結果から、これらの人々においては、社会環境が不良なためではなく、鉛の曝露による長期の神経学的影響がその後の認知機能および行動上の問題の重要な原因であるとの過去の研究を支持するものである。

Full Text

A study using functional magnetic resonance imaging (fMRI) to evaluate brain function revealed that adults who were exposed to lead as children incur permanent brain injury. The results were presented at the annual meeting of the Radiological Society of North America (RSNA).

"What we have found is that no region of the brain is spared from lead exposure," said the study's lead author, Kim Cecil, Ph.D., imaging scientist at Cincinnati Children's Hospital Medical Center and professor of radiology, pediatrics and neuroscience at the University of Cincinnati College of Medicine. "Distinct areas of the brain are affected differently."

The study is part of a large research project called the Cincinnati Lead Study, a long-term lead exposure study conducted through the Cincinnati Children's Environmental Health Center, a collaborative research group funded by the National Institute of Environmental Health Sciences and U.S. Environmental Protection Agency. The Cincinnati Lead Study followed prenatal and early childhood lead exposure of 376 infants from high-risk areas of Cincinnati between 1979 and 1987. Over the course of the project, the children underwent behavioral testing and 23 blood analyses that yielded a mean blood lead level.

Lead, a common and potent poison found in water, soil and lead-based paint, is especially toxic to children's rapidly developing nervous systems. Homes built before 1950 are most likely to contain lead-based paint, which can chip and be ingested by children.

"Lead exposure has been associated with diminished IQ, poor academic performance, inability to focus and increased risk of criminal behavior," Dr. Cecil said.

Dr. Cecil's study involved 33 adults who were enrolled as infants in the Cincinnati Lead Study. The mean age of the study participants, which included 14 women and 19 men, was 21 years. The participants' mean blood lead levels ranged from 5 to 37 micrograms per deciliter with a mean of 14. Participant histories showed IQ deficiencies, juvenile delinquency and a number of criminal arrests.

Each participant underwent fMRI while performing two tasks to measure the brain's executive functioning, which governs attention, decision-making and impulse control. The imaging revealed that in order to complete a task that required inhibition, those with increased blood lead levels required activation from additional regions within the frontal and parietal lobes of the brain.

"This tells us that the area of the brain responsible for inhibition is damaged by lead exposure and that other regions of the brain must compensate in order for an individual to perform," Dr. Cecil said. "However, the compensation is not sufficient."

Imaging performed during a second task designed to test attention revealed an association between higher lead levels and decreased activation in the parietal region and other areas of the brain.

According to Dr. Cecil, the brain's white matter, which organizes and matures at an early age, adapts to lead exposure, while the frontal lobe, which is the last part of the brain to develop, incurs multiple insults from lead exposure as it matures.

"Many people think that once lead blood levels decrease, the effects should be reversible, but, in fact, lead exposure has harmful and lasting effects," she said.

Dr. Cecil believes that these findings lend support to previous reports from the Cincinnati Lead Study showing that the lasting neurological effect of lead exposure, rather than a poor social environment, is a key contributor to the subsequent cognitive and behavior problems in this group.

Co-authors are Kim M. Dietrich, Ph.D., M.S., Caleb M. Adler, M.D., James C. Eliassen, Ph.D., and Bruce P. Lanphear, M.D., M.P.H.

TOPICS

Cardiology

心疾患の基礎にあるPADの徴候

冠動脈CT血管造影は胸痛のトリアージに役立つ

Oncology

エラストグラフィーにより不必要な乳房生検が減少する

乳房生検の代替策としての超音波検査

エラストグラフィーは皮膚がんを正確に見極める

年1回の乳房超音波検査は一部の女性においては有益である

Psychiatry

鉛への曝露は長期に渡り認知機能に対し影響する

情動刺激の処理過程には性差がある